

10th Science Lesson 12 Notes in English

12] Plant Anatomy and Plant Physiology

Introduction

- Plants exhibit varying degrees of organization. Atoms are organized into molecules, molecules into organelles, organelles into cells, cells into tissues and tissues into organs.
- The first account of internal structure of plants was published by English Physician
- **Nehemiah Grew**. He is known as **Father of Plant Anatomy**. Plant anatomy (Gk *Ana* = as under; *Temnein* = to cut) is the study of internal structure of plants.
- You have already studied the different kinds of tissues in standard IX. In this lesson, you will study about the internal structure of plant tissues, process of photosynthesis and respiration.

Tissues

- Tissues are the group of cells that are similar or dissimilar in structure and origin, but perform similar function.
- Plant tissues Tissues are the group of cells that are similar or dissimilar in structure and origin, but perform similar function.
- Plant tissues can be broadly classified into two, based on their ability to divide. They are
 - i) Meristamatic tissue
 - ii) Permanent tissue.

Tissue system

Sachs (1875) classified tissue system in plants into three types

- i) Dermal or Epidermal tissue system
- ii) Ground tissue system
- iii) Vascular tissue system

Dermal or Epidermal Tissue System

- It consists of epidermis, stomata and epidermal outgrowths.
- Epidermis is the outer most layer.
- It has many minute pores called stomata.
- Cuticle is present on the outer wall of epidermis to check evaporation of water.
- Trichomes and root hairs are the epidermal outgrowths.

Tissue system and its function

Tissue System	Components	Functions
Dermal Tissue System	Epidermis and Periderm (in older stems and roots)	<ul style="list-style-type: none"> • Protection • Prevention of water loss
Ground Tissue System	Parenchyma tissue Collenchyma tissue Sclerenchyma tissue	<ul style="list-style-type: none"> • Photosynthesis • Food storage • Regeneration • Support • Protection
Vascular Tissue System	Vascular tissues - Xylem tissue - Phloem tissue	<ul style="list-style-type: none"> • Transport of water and minerals • Transport of food

Functions:

- Epidermis protects the inner tissues.
- Stomata helps in transpiration.
- Root hairs help in absorption of water and minerals.

Ground Tissue System

- It includes all the tissues of the plant body except epidermal and vascular tissues like (i) Cortex (ii) Endodermis (iii) Pericycle (iv) Pith

Vascular Tissue System

- It consists of xylem and phloem tissues.
- They are present in the form of bundles called vascular bundles. Xylem conducts water and minerals to different parts of the plant. Phloem conducts food materials to different parts of the plant.
- There are three different types of vascular bundles namely (i) Radial (ii) Conjoint (iii) Concentric

(i) Radial Bundles

- Xylem and phloem are present in different radii alternating with each other. e.g. roots

ii) Conjoint bundles

- Xylem and phloem lie on the same radius. There are two types of conjoint bundles.

a) Collateral

- Xylem lies towards the centre and phloem lies towards the periphery.
- When cambium is present in collateral bundles, it is called open. e.g. dicot stem and collateral bundle without cambium is called closed. e.g. monocot stem.

b) Bicollateral

- In this type of bundle, the phloem is present on both outer and inner side of xylem. e.g. Cucurbita

(iii) Concentric Bundles

- Vascular bundle in which xylem completely surrounds the phloem or viceversa is called concentric vascular bundle. It is of two types:

1. Amphivasal:

- Xylem surrounds phloem. e.g. *Dracaena*

2. Amphicribal:


- Phloem surrounds xylem. e.g. Ferns

Endarch:

- Protoxylem lies towards the centre and metaxylem lies towards the periphery. e.g. stem.

Exarch :

- Protoxylem lies towards the periphery and metaxylem lies towards the centre. e.g. roots.



Types of vascular bundle

Internal Structure of Dicot Root (Bean)

A thin transverse section of dicot root shows the following structures.

(i) Epiblema:

- It is the outermost layer.
- Cuticle and stomata are absent. Unicellular root hairs are present. It is also known as **Rhizodermis** or **Piliferous layer**.

(ii) Cortex:

- It is a multilayered large zone made of thin-walled parenchymatous cells with intercellular spaces.
- It stores food and water.

(iii) Endodermis:

- It is the innermost layer of cortex.
- The cells are barrel - shaped, closely packed, and show band like thickenings on their radial and inner tangential walls called **casparian strips**.

- But these casparian strips are absent in the endodermis cells which are located opposite the protoxylem these thin walled cells without casparian strips are called passage cell.
- It helps in the movement of water and dissolved salts from cortex into xylem.

(iv) Stele:

- All tissues inner to endodermis constitute stele.
- It includes pericycle and vascular bundle.

(a) Pericycle:

- Inner to endodermis lies a single layer of pericycle. It is the site of origin of lateral roots.

(b) Vascular bundle:

- It is radial.
- Xylem is **exarch** and **tetrach**. The tissue present between xylem and phloem is called conjunctive tissue. In dicot root, it is made up of parenchyma.

(c) Pith:


- Young root contains pith whereas in old root pith is absent.

Internal Structure of Monocot Root (Maize)

- A thin transverse section of monocot root, shows the following characteristic features.

i. Epiblema or Rhizodermis:

- It is the outermost layer of the root, and is made up of single layer of thin walled, parenchymatous cell.
- Stomata and cuticle are absent. The root hair helps in absorption of water and minerals from the soil.
- This layer also protects the inner tissues.



Transverse section of Dicot root

ii. Cortex:

- It is multilayered large zone, composed of parenchymatous cells with intercellular spaces. It stores water and food material.

iii. Endodermis:

- It is the innermost layer of cortex with characteristic casparian strips and passage cells. **Casparian strips** are band like thickening made of **suberin**.

iv. Stele:



- All the tissues inner to endodermis constitute stele. It includes pericycle, vascular tissues and pith.

a) Pericycle:

- It is a single layer of thin walled cells. The lateral roots originate from this layer.

b) Vascular tissues:

- It consists of many patches of xylem and phloem arranged radially. The xylem is exarch and polyarch.
- The conjunctive tissue is made up of sclerenchyma.



Transverse section of Monocot root

c) Pith:

- It is present at the center. It is made up of parenchyma cells with intercellular spaces. It contains abundant amount of starch grains. It stores food.

Internal Structure of Dicot Stem (Sunflower)

The transverse section of a dicot stem reveals the following structures.

1. Epidermis:

- It is the outermost layer. It is made up of single layer of parenchyma cells, its outer wall is covered with cuticle. It is protective in function.

2. Cortex:-

- It is divided into three regions:

(i) Hypodermis:

- It consists of 3 - 6 layers of collenchyma cells. It gives mechanical support.

Differences between Dicot and Monocot root

S. No.	Tissues	Dicot Root	Monocot Root
1	Number of Xylem	Tetrarch	Polyarch
2	Cambium	Present (During secondary growth only)	Absent
3	Secondary Growth	Present	Absent
4	Pith	Absent	Present
5	Conjunctive Tissue Ex.	Parenchyma Bean	Sclerenchyma Maize

(ii) Middle cortex:

- It is made up of few layers of chlorenchyma cells. It is involed in photosynthesis due to the presence of chloroplast.

(iii) Inner cortex:

- It is made up of few layers of parenchyma cells. It helps in gaseous exchange and stores food materials.
- **Endodermis** is the inner most layer of cortex it consists of a single layer of barrel shaped cells, these cells contain starch grains. So it is also called **starch sheath**.

3. Stele:

- The central part of the stem inner to endodermis is known as stele. It consists of pericycle, vascular bundle and pith.

(i) Pericycle:

- It occurs between vascular bundle and endodermis. It is multilayered, parenchymatous with alternating patches of sclerenchyma.

(ii) Bundle Cap:

- There is a patch of hard sclerenchyma tissue outside to the phloem of vascular bundle is called Bundle Cap.

(iii) Vascular bundle:

- Vascular bundles are conjoint, collateral, endarch and open. They are arranged in the form of a ring around the pith.

(iv) Pith:

- The large central parenchymatous zone with intercellular spaces is called pith. It helps in the storage of food materials.

Internal Structure of Monocot Stem (Maize)

A transverse section of monocot stem reveals the following structures.

1. Epidermis:

- It is the outermost layer. It is made up of single layer of parenchyma cells. It is covered with thick cuticle. Multicellular hairs are absent and stomata are also less in number.

2. Hypodermis:

- It is made up of few layers of sclerenchyma cells interrupted by chlorenchyma. Sclerenchyma provides mechanical support to plant.

3. Ground tissue:

- The entire mass of parenchyma cells next to hypodermis and extending to the centre is called **ground tissue**. It is not differentiated into endodermis, cortex, pericycle and pith.

4. Vascular Bundle:

- Vascular bundles are skull shaped and scattered in the ground tissue.
- Vascular bundles are conjoint, collateral, endarch and closed. Each vascular bundle is surrounded by few layer of sclerenchyma cells called **bundle sheath**.

(a) Xylem:



- It consists of metaxylem and protoxylem. Xylem vessels are arranged in V or Y shape. In mature vascular bundle, the lower most protoxylem disintegrates and form a cavity. This is called **protoxylem lacuna**.

(b) Phloem:

- It consists of sieve tube elements and companion cells. Phloem parenchyma, and phloem fibers are absent.

5. Pith:

- Pith is not differentiated in monocot stems.



Transverse section of Dicot stem

Differences between Dicot Stem Ex. Sunflower and Monocot Stem Ex. Maize


S. No.	Tissues	Dicot Stem	Monocot Stem
1	Hypodermis	Collenchymatous	Sclerenchymatous
2	Ground tissue	Differentiated into cortex, endodermis, pericycle and pith	Undifferentiated
3	Vascular bundles	(i) Less in number (ii) Uniform in size (iii) Arranged in a ring (iv) Open (Cambium present) (v) Bundle sheath absent	(i) Numerous (ii) Smaller near periphery, bigger in the centre (iii) Scattered (iv) Closed (Cambium absent) (v) Bundle sheath present
4	Secondary growth	Present	Mostly absent
5	Pith	Present	Absent
6	Medullary rays	Present	Absent

Internal Structure of Dicot Leaf (Dorsiventral Leaf) Ex. Mango Leaf

The transverse section of leaf shows the following structures.

(i) Upper epidermis:

- This is the outermost layer made of single layered parenchymatous cells without intercellular spaces. The outer wall of the cells are cuticularized. Stomata are less in number.



(ii) Lower epidermis:

- It is a single layer of parenchymatous cells with a thin cuticle. It contains numerous stomata. Chloroplasts are present only in guard cells.
- The lower epidermis helps in the exchange of gases.
- The loss of water vapour is facilitated through this chamber.

(iii) Mesophyll:

- The tissue present between the upper and lower epidermis is called mesophyll. It is differentiated into Palisade parenchyma and Spongy parenchyma.

a) Palisade parenchyma:

- It is found just below the upper epidermis. The cells are elongated.


- These cells have more number of chloroplasts. The cells do not have intercellular spaces and they take part in photosynthesis.

b) Spongy parenchyma:

- It is found below the palisade parenchyma tissue. Cells are almost spherical or oval and are irregularly arranged. Cells have intercellular spaces. It helps in gaseous exchange.

(iv) Vascular bundles:

- Vascular bundle is present in mid-rib and lateral veins. Vascular bundles are conjoint, collateral and closed.
- Each vascular bundle is surrounded by a sheath of parenchymatous cells called **bundle sheath**. Each vascular bundle consists of xylem lying towards the upper epidermis and phloem towards the lower epidermis.



Transverse section of Dicot leaf

Internal Structure of Monocot Leaf (Isobilateral Leaf) Ex. Grass Leaf

The transverse section of a monocot leaf reveals the following structures.

(i) Epidermis:

- Monocot leaf has upper and lower epidermis. Epidermis is made up of parenchyma cells. Cuticle is present on the outer wall stomata are present on both upper and lower epidermis.
- Some cells of upper epidermis are large and thin walled they are known as **bulliform cells**.


(ii) Mesophyll:

- It is the ground tissue that is present between both epidermal layers.

- Mesophyll is not differentiated into palisade and spongy parenchyma. The cells are irregularly arranged with inter-cellular spaces. These cells contain chloroplasts.

(iii) Vascular bundles:

- Large number of vascular bundles are present, some of which are small and some are large.
- Each vascular bundle is surrounded by parenchymatous bundle sheath. Vascular bundles are conjoint, collateral and closed.
- Xylem is present towards upper epidermis and phloem towards lower epidermis.



Transverse section of Monocot Leaf

Differences between of Dicot and Monocot Leaf

S. No.	Dicot Leaf	Monocot Leaf
1	Dorsiventral leaf	Isobilateral leaf
2	Mesophyll is differentiated into palisade and spongy parenchyma	Mesophyll is not differentiated into palisade and spongy parenchyma

Plant Physiology

Plastids

- Plastids are double membrane bound organelles found in plants and some algae. They are responsible for preparation and storage of food. There are three types of plastids.

Chloroplast - green coloured plastids

Chromoplast - yellow, red, orange coloured plastids

Leucoplast - colourless plastids

Structure of Chloroplast

- Chloroplasts are green plastids containing green pigment called **chlorophyll**. Chloroplasts are oval shaped organelles having a diameter of 2-10 micrometer and a thickness of 1-2 micrometer.



Ultra structure of Chloroplast

1. Envelope:

- Chloroplast envelope has outer and inner membranes which is separated by inter membrane space.

2. Stroma:

- Matrix present inside to the membrane is called stroma. It contains DNA, 70 S ribosomes and other molecules required for protein synthesis.

3. Thylakoids:

- It consists of thylakoid membrane that encloses thylakoid lumen.
- Photosynthetic pigments are present in thylakoids. Thylakoids forms a stack of **disc like structures** called a grana (singular granum).

4. Grana:


- Thylakoids arranged in the form of discs stacked one above the other called granum. Grana are inter connected by stroma lamella.

Functions of Chloroplast

1. Photosynthesis 2. Storage of starch 3. Synthesis of fatty acids 4. Storage of lipids 5. Formation of chloroplasts

Photosynthesis

- Photosynthesis (Photo = light; synthesis = to build) is a process by which autotrophic organisms like green plants, algae and chlorophyll containing bacteria utilize the energy from sunlight to synthesize their own food.
- In this process, carbon dioxide combines with water in the presence of sunlight and chlorophyll to form carbohydrates. During this process oxygen is released as a byproduct.



Carbon dioxide + Water \longrightarrow Glucose + Water + Oxygen

Where does photosynthesis occur?

- Photosynthesis occurs in all green parts of the plant especially in green leaves.

Photosynthetic Pigments

- Pigments involved in photosynthesis are called **Photosynthetic pigments**.
- Photosynthetic pigments are of two classes namely, the primary pigments and accessory pigments.
- Chlorophyll a is the **primary pigment** that traps solar energy and converts it into electrical and chemical energy.
- Thus it is called the reaction centre. Other pigments such as chlorophyll b and carotenoids are called **accessory pigments** as they pass on the absorbed energy to chlorophyll a (Chl.a) molecule. Reaction centre (Primary pigments) and harvesting centre (Accessory pigments) together form Pigment systems.

Role of Sunlight in Photosynthesis

- The entire process of photosynthesis takes place inside the chloroplast.
- The structure of chloroplast is such that the light dependent (**Light reaction**) and light independent (**Dark reaction**) take place at different sites in the organelle


1. Light dependent Reaction (Hill reaction \ Light reaction)

- This was discovered by **Robin Hill** (1939). This reaction takes place in the presence of light energy in **thylakoid membranes** (grana) of the chloroplasts.

- Photosynthetic pigments absorb the light energy and convert it into chemical energy ATP and NADPH₂. These products of light reaction move out from the thylakoid to the stroma of the chloroplast.

2. Light independent reactions (Dark reaction) (Biosynthetic phase)

- Dark reaction or biosynthetic pathway is takes place in **stroma**. During this reaction CO₂ is reduced into carbohydrates with the help of light generated **ATP** and **NADPH₂**.
- This is also called as **Calvin cycle** and is carried out in the absence of light. It is called dark reaction.
- In Calvin cycle the inputs are CO₂ from the atmosphere and the ATP and NADPH₂ produced from light reaction.



Overview of Hill and Calvin cycle

Factors Affecting Photosynthesis

a) Internal Factors:

- i) Pigments ii) Leaf age iii) Accumulation of carbohydrates iv) Hormones

b) External Factors:

- i) Light ii) Carbon dioxide iii) Temperature iv) Water v) Mineral elements


Mitochondria

- Mitochondria are filamentous or granular cytoplasmic organelles present in cells.
- The mitochondria were first discovered by Kolliker in 1857 as granular structures in striated muscles.

- Mitochondria (singular: mitochondrion) are organelles within eukaryotic cells that produce adenosine triphosphate (ATP) which form the energy currency of the cell, for this reason, the mitochondria is referred to as the **"Power house of the cell"**. Mitochondria vary in size from 0.5 μm to 2.0 μm .
- Mitochondria contain 60-70% protein, 25-30% lipids, 5-7% RNA and small amount of DNA and minerals.

Structure of Mitochondria Mitochondrial Membranes:

- It consists two membranes called inner and outer membrane. Each membrane is 60-70 \AA thick.
- Outer mitochondrial membrane is smooth and freely permeable to most small molecules.
- It contains enzymes, proteins and lipids. It has **porin molecules** (proteins) which form channels for passage of molecules through it.



Structure of Mitochondria


- Inner mitochondrial membrane is semi permeable membrane and regulates the passage of materials into and out of the mitochondria.
- It is rich in enzymes and carrier proteins. It consists of 80% proteins and lipids.

Cristae:

- The inner mitochondrial membrane gives rise to finger like projections called cristae. These cristae increase the inner surface area (fold in inner membrane) of the mitochondria to hold variety of enzymes.

Oxysomes:

- The inner mitochondrial membrane bear minute regularly spaced tennis racket shaped particles known as oxysomes (F₁ particle). They involve in ATP synthesis.



Structure of Oxysomes

- Mitochondrial matrix** - It is a complex mixture of proteins and lipids. Matrix contains enzymes for Krebs cycle, mitochondrial ribosomes (70S), tRNAs and mitochondrial DNA.

Functions of Mitochondria

- Mitochondria is the main organelle of cell respiration. They produce a large number of ATP molecules. So they are called as **power houses of the cell** or **ATP factory of the cell**.
- It helps the cells to maintain normal concentration of calcium ions.
- It regulates the metabolic activity of the cell.

Types of Respiration

- Respiration involves exchange of gases between the organism and the external environment.
- The plants obtain oxygen from their environment and release carbon dioxide and water vapour. This exchange of gases is known as **external respiration**. It is a physical process.
- Biochemical process occurs within cells where the food is oxidized to obtain energy, this is known as cellular respiration.



Stages of Aerobic respiration

a. Glycolysis (Glucose splitting):

- It is the breakdown of one molecule of glucose (6 carbon) into two molecules of pyruvic acid (3 carbon). Glycolysis takes place in cytoplasm of the cell. It is the first step of both aerobic and anerobic respiration.

b. Krebs Cycle:

- This cycle occurs in mitochondria matrix. At the end of glycolysis, 2 molecules of pyruvic acid enter into mitochondria.
- The oxidation of pyruvic acid into CO₂ and water takes place through this cycle. It is also called **Tricarboxylic Acid Cycle** (TCA).

c. Electron Transport Chain:

- This is accomplished through a system of electron carrier complex called **electron transport chain** (ETC) located on the inner membrane of the mitochondria.
- NADH₂ and FADH₂ molecules formed during glycolysis and Krebs cycle are oxidised to NAD⁺ and FAD⁺ to release the energy via electrons.
- The electrons, as they move through the system, release energy which is trapped by ADP to synthesize ATP. This is called **oxidative phosphorylation**. In this process, O₂ the ultimate acceptor of electrons gets reduced to water.

Anaerobic respiration

- Anaerobic respiration takes place without oxygen. Glucose is converted into ethanol (Ethanol fermentation by yeast) or lactic acid (lactic acid fermentation by bacteria).

Respiratory quotient (R.Q)

- Respiratory quotient is the ratio of volume of carbon dioxide liberated and the volume of oxygen consumed during respiration. It is expressed as

$$RQ = \frac{\text{Volume of CO}_2 \text{ liberated}}{\text{Volume of O}_2 \text{ consumed}}$$

More to know:

ATP	Adenosine Triphosphate
ADP	Adenosine Diphosphate
NAD	Nicotinamide Adenine Dinucleotide
NADP	Nicotinamide Adenine Dinucleotide Phosphate

- A cell cannot get its energy directly from glucose. So in respiration the energy released from glucose is used to make ATP (Adenosine Triphosphate)
- Melvin Calvin, an American biochemist, discovered chemical pathway for photosynthesis. The cycle is named as Calvin cycle. He was awarded with Nobel Prize in the year 1961 for his discovery.
- Artificial photosynthesis is a method for producing renewable energy by the use of sunlight. Indian scientist C.N.R. Rao who was conferred the Bharat Ratna (2013) is also working on similar technology of artificial photosynthesis to produce -Hydrogen fuel (renewable energy).